



ANTI-DEAMIDATED GLIADIN PEPTIDE (DGP) IgG

Synonym(s): DGP

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

Lab Section Category	Immunology/Serology/STD/Allergy - Immunology Click here to find out more about the write-up (/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/immunologyserologystdallergy.aspx).
Indications	Anti-tTG IgG and anti-DGP IgG should be performed in patients in whom low IgA or selective IgA deficiency is identified. Anti-DGP (IgA and IgG) should be combined with the anti-tTG IgG test when screening children <2 years of age for Celiac disease.
Specimen Required	3 mL plain blood
Method	ELISA Immunoassay
Test Result	Reported numerically
Reference Interval/Value	<10 U/mL Negative > =10 U/mL Positive
Turnaround Time	Range 2 to 3 working days
Day(s) Test Set up	Mondays to Fridays
Remarks	Diagnostic testing should be done with patients on gluten-containing diets.

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

Change History Notes

- **29 Feb 2016 01:57 PM**
New test available w.e.f 1 March 2016

©Copyright 2018 Singapore General Hospital, Outram Road, Singapore 169608