



ANTI-MPO AND ANTI-PR3 ANTIBODIES (ANCAE)

Synonym(s): Anti MPO, MPO Antibody, Anti Myeloperoxidase, Anti PR3, PR3 Antibody, Anti Proteinase 3

Back To Previous Page (<http://cms-sgh.sppub/Clinical-Departments-Centers/Pathology/Pathology-Handbook/Lab-Discipline-Special-Instructions/Tests/Pages/Forms/AllItems.asp?category=SW1tdW5vbG9neS9TZXJvbG9neS9TVEQvQWxsZXJneSAtIEltbXVub2xvZ3k%3d-qLAjr1Ja5yM%3d>) (<http://cms-sgh.sppub/Clinical-Departments-Centers/Pathology/Pathology-Handbook/Lab-Discipline-Special-Instructions/Tests/Pages/Forms/AllItems.aspx?category=SW1tdW5vbG9neS9TZXJvbG9neS9TVEQvQWxsZXJneSAtIEltbXVub2xvZ3k%3d-qLAjr1Ja5yM%3d>)

Lab Section Category	Immunology/Serology/STD/Allergy - Immunology Click here to find out more about the write-up (/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/immunologyserologystdallergy.aspx).
Specimen Required	3 mL plain blood
Method	Enzyme Immunoassay
Test Result	Reported numerically
Reference Interval/Value	<ul style="list-style-type: none"> – Anti-MPO <ul style="list-style-type: none"> < 20 RU/mL Negative >= 20 RU/mL Positive – Anti-PR3 <ul style="list-style-type: none"> < 20 RU/mL Negative >= 20 RU/mL Positive
Turnaround Time	Range 2 – 4 days
Day(s) Test Set up	Batch tested, Monday, Wednesday and Friday

Back To Previous Page (<http://cms-sgh.sppub/Clinical-Departments-Centers/Pathology/Pathology-Handbook/Lab-Discipline-Special-Instructions/Tests/Pages/Forms/AllItems.asp?category=SW1tdW5vbG9neS9TZXJvbG9neS9TVEQvQWxsZXJneSAtIEltbXVub2xvZ3k%3d-qLAjr1Ja5yM%3d>) (<http://cms-sgh.sppub/Clinical-Departments-Centers/Pathology/Pathology-Handbook/Lab-Discipline-Special-Instructions/Tests/Pages/Forms/AllItems.aspx?category=SW1tdW5vbG9neS9TZXJvbG9neS9TVEQvQWxsZXJneSAtIEltbXVub2xvZ3k%3d-qLAjr1Ja5yM%3d>)

