



## BETA-CROSSLAPS, PLASMA

### Synonym(s):

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

<b>Lab Section Category</b>	<b>Clinical Biochemistry</b> Click here to find out more about the <b>write-up</b> ( <a href="/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/clinical-biochemistry.aspx">/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/clinical-biochemistry.aspx</a> ).
<b>Specimen Required</b>	3 mL EDTA blood  Preferably fasting morning sample. Send sample immediately to the laboratory.
<b>Method</b>	Immunoassay
<b>Reference Interval/Value</b>	>12 yrs: M 0.16 - 0.88 µg/L ; F (≤50 yrs) 0.10 - 0.78 µg/L ; F (≥51 yrs) 0.06 - 1.10 µg/L
<b>Turnaround Time</b>	Mean = 4 hrs, Range: 3 – 5 hrs
<b>Day(s) Test Set up</b>	Monday – Saturday

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

©Copyright 2018 Singapore General Hospital, Outram Road, Singapore 169608

