



	Make/Change Appointment		Pay Online
---	---	---	----------------------------

FACTOR ASSAYS

Synonym(s): Fibrinogen Assay, Factor I Assay, Factor II Assay

Back To Previous Page ([javascript: history.go\(-1\)](javascript: history.go(-1);))

Lab Section Category	Haematology Click here to find out more about the write-up (/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/haematology.aspx).
Specimen Required	3 mL blood containing 0.3 mL of sodium citrate Patient must not be on heparin.
Method	Clotting Assay

Reference Interval/Value	Factor assay	Reference range
	I(Fibrinogen)	1.8 – 4.8 g/L
	II(Prothrombin)	80 – 160%
	V	70 – 170%
	VII	40 – 180%
	VIII	55 – 200%
	IX	40 – 200%
	X	60 – 160%
	XI	70 – 200%
	XII	30 – 200%
Turnaround Time	2 days	
Day(s) Test Set up	Monday - Saturday	
Remarks	Specific Coagulation factor assays are available individually. Each assay utilises the same sample requirement. Order the specific factor required.	

[Back To Previous Page \(javascript: history.go\(-1\)\)](javascript: history.go(-1))

©Copyright 2018 Singapore General Hospital, Outram Road, Singapore 169608