SharePoint Newsfeed (http://mysite.sppub:80/default.aspx) OneDrive (http://mysite.sppub:80/\_layouts/15/MySite.aspx?MySiteRedirect=AllDocuments) Sites (h





## IDENTIFICATION OF MYCOBACTERIA

**Synonym(s):** Mycobacterial Identification

Back To Previous Page (javascript: history.go(-1))

Lab Section Category	Mycobacteriology Click here to find out more about the write-up (/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/mycobacteriology.aspx).
Specimen Required	Referred cultured acid-fast positive isolate from other laboratories
Storage and Transport	Fresh pure culture of isolate on slant in sealed, screw-capped plastic tube. Local laboratories may submit positive MGIT tubes or Myco-F lytic cultures, but these must be packed and labelled properly.
	All containers should be tightly closed, labeled with identifying lab number and placed in a biohazard-labelled ziplock bag. Pack and transport according to MOH-BATA requirements.
	Request form (stating name and address of referring laboratory, source of mycobacterial isolate, patient particulars with relevant clinical information, clinician in-charge and requester's contact number) is to be placed in separate sealed plastic bag outside the ziplock bag. The type of test should be specified as <b>Mycobacterial Identification</b> . For regional labs, packaging must conform to IATA standards.
Method	AFB smear, culture, DNA probes, line probe assays and MALDI-TOF.
Test Result	Species identified reported
Turnaround Time	2 - 4 weeks depending on the nature of the organism
Day(s) Test Set up	Monday - Friday
<	>

Remarks	Please contact CTBL at tel: 6576 7655/7656 before sending isolate	
Back To Previous Page (javascript: history.go(-1))  Change History Notes		
29 Apr 2016 05:26 PM     Amend Storage and Transport.		
12 Jan 2018 11:58 AM     Remove HPLC in methods		
©Copyright 2018 Singapore General Hospital, Out	tram Road, Singapore 169608	