

SharePoint [Newsfeed \(http://mysite.sppub:80/default.aspx\)](http://mysite.sppub:80/default.aspx) [OneDrive \(http://mysite.sppub:80/_layouts/15/MySite.aspx?MySiteRedirect=AllDocu\)](http://mysite.sppub:80/_layouts/15/MySite.aspx?MySiteRedirect=AllDocu)



	Make/Change Appointment		Pay Online
---	---	---	----------------------------

IMMUNOFIXATION ELECTROPHORESIS, URINE

Synonym(s):

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

Lab Section Category	Clinical Biochemistry Click here to find out more about the write-up (/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/clinical-biochemistry.aspx) .
Specimen Required	20 mL fresh random urine in sterile container.
Method	Agarose Gel Electrophoresis / Immunodetection
Reference Interval/Value	No monoclonal bands detected (kappa, lambda light chains)
Turnaround Time	Mean = 7 days, Range: 3 – 11 days
Day(s) Test Set up	Monday - Friday

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

©Copyright 2018 Singapore General Hospital, Outram Road, Singapore 169608

