



## MICROALBUMIN, RANDOM URINE

**Synonym(s):**

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

<b>Lab Section Category</b>	<b>Clinical Biochemistry</b> Click here to find out more about the <b>write-up</b> ( <a href="/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/clinical-biochemistry.aspx">/clinical-departments-centers/pathology/pathology-handbook/lab-discipline-special-instructions/pages/clinical-biochemistry.aspx</a> ).
<b>Specimen Required</b>	20 mL fresh random urine in sterile container.
<b>Method</b>	Immunoassay
<b>Reference Interval/Value</b>	>12 yrs : <19 mg/L
<b>Turnaround Time</b>	Mean = 4 hrs, Range: 3 – 5 hrs
<b>Day(s) Test Set up</b>	Monday – Saturday
<b>Remarks</b>	<p>Instructions before sampling:</p> <ol style="list-style-type: none"> <li>1. Patient should be well as baseline. They should have no urinary tract infection, no acute febrile illness, no intense exercise within the previous 24 hrs and not be menstruating.</li> <li>2. Recommended urine collection is a fresh, first morning void sample. If a first morning void is not practical, random spot samples are acceptable.</li> </ol>

[Back To Previous Page \(javascript: history.go\(-1\)\)](#)

©Copyright 2018 Singapore General Hospital, Outram Road, Singapore 169608

